arXiv:1601.03954v2 [math.NT] 25 May 2016

QUANTUM j-INVARIANT IN POSITIVE CHARACTERISTIC II: FORMULAS AND VALUES AT THE QUADRATICS

L. DEMANGOS AND T.M. GENDRON

Abstract. In this sequel to [2], the multi-valued quantum j-invariant in positive characteristic is studied at quadratic elements. For every quadratic f, an explicit expression for each of the values of $j^{\mathrm{qt}}(f)$ is given as a limit of rational functions of f. It is proved that the number of values of $j^{\mathrm{qt}}(f)$ is finite.

1. Introduction

Let $\mathbb{F}_q$ be the field with q elements, $q = p^r$ a prime power, $k = \mathbb{F}_q(T)$ the function field over $\mathbb{F}_q$, k_∞ the completion of k with respect to the valuation $v(f) = -\deg_T(f)$. In [2], the quantum j-invariant was introduced as a multi-valued modular function

$$j^{\mathrm{qt}} : \mathrm{GL}_2(A)\backslash k_\infty \multimap k_\infty \cup \{\infty\}$$

obtained as the limit of a sequence of approximating functions

$$j_\varepsilon : k_\infty \longrightarrow k_\infty \cup \{\infty\} \tag{1}$$

as $\varepsilon \to 0$. Explicit formulas for the approximants (1) in terms of the sequence $\{\mathfrak{q}_i\} \subset A$ of best approximations of f were obtained and using them, it was shown that $f \in k$ if and only if eventually $j_\varepsilon(f) = \infty$.

In this paper we consider $j^{\mathrm{qt}}(f)$ in the special case of $f \in k_\infty$ quadratic. Here we are able to derive explicit formulas not just for the approximants but for all the values of $j^{\mathrm{qt}}(f)$, each of which expressed as a limit of certain rational functions of f. Using these formulas we are able to prove that for f quadratic, $\#j^{\mathrm{qt}}(f) < \infty$.

In what follows, we fix notation used in [2].

2. Values at Quadratic Units

Fix $a \in A - \mathbb{F}_q$, $b \in \mathbb{F}_q^\times$ with

$$d := \deg a > 0$$

and write $D = a^2 + 4b$. The solutions f, f' of $X^2 - aX - b = 0$ are quadratic units, and every quadratic unit in $k_\infty - k$ occurs in this way. Since $ff' = -b$, we have $|ff'| = 1$. Moreover, since $f + f' = a$, we have as well that $|f|, |f'| \neq 1$. Therefore, if we assume that $|f| > 1$, then

$$|f| = |a| = q^d > 1 \quad \text{and} \quad |f'| = |b/a| = q^{-d} < 1.$$

Consider the recursive sequence

$$\mathtt{Q}_0 = 1, \mathtt{Q}_1 = a, \ \ldots, \ \mathtt{Q}_{n+1} = a\mathtt{Q}_n + b\mathtt{Q}_{n-1}.$$

1991 *Mathematics Subject Classification.* Primary 11R58,11F03,11R11; Secondary 11K60.

Key words and phrases. quantum j-invariant, global function fields, quadratic extensions, Diophantine approximation.

Note that when $b = 1$, $\mathtt{Q}_n = \mathtt{q}_n$ is the sequence of best approximations for $f = [a, a, \dots]$; see §3 of [2]. It is clear that for all n

$$|\mathtt{Q}_n| = |a|^n = q^{dn}.$$

Recall *Binet's formula*

$$\mathtt{Q}_n = \frac{f^{n+1} - (f')^{n+1}}{\sqrt{D}}, \quad n = 0, 1, \dots.$$

which, while usually stated over $\mathbb{Q}$, is equally true in this setting, by a very simple induction on n. The root $\sqrt{D}$ is chosen so that in odd characteristic, $f = (a + \sqrt{D})/2$; in even characteristic, we simply take $\sqrt{D} = a$.

Recall that $\|x\|$ = distance of x to the nearest element of A; see [2], §2 for this and other relevant notation. Binet's formula immediately gives

$$\|\mathtt{Q}_n f\| = q^{-(n+1)d} : \tag{2}$$

indeed,

$$|\mathtt{Q}_n f - \mathtt{Q}_{n+1}| = \frac{|f'|^{n+1}|f - f'|}{|\sqrt{D}|} = |f'|^{n+1} = q^{-(n+1)d} < 1. \tag{3}$$

In particular, $\mathtt{Q}_n^{\perp} = \mathtt{Q}_{n+1}$, where $\lambda^{\perp}$ denotes the element of A closest to λf.

The remainder of this section is devoted to deriving an explicit formula for each value of $j^{\mathrm{qt}}(f)$. In view of our need to work with monic polynomials, we make the following modification to the above sequence. For each n, we write $\bar{\mathtt{Q}}_n$ for the unique monic polynomial obtained as $c_n \mathtt{Q}_n$ for some $c_n \in \mathbb{F}_q^{\times}$. Then $|\bar{\mathtt{Q}}_n| = |\mathtt{Q}_n|$ and $\|\bar{\mathtt{Q}}_n f\| = \|\mathtt{Q}_n f\|$ since $\bar{\mathtt{Q}}_n^{\perp} = c_n \mathtt{Q}_n^{\perp}$. (N.B. $\bar{\mathtt{Q}}_n^{\perp}$ is not necessarily monic.) Note that if we choose $c \in \mathbb{F}_q^{\times}$ so that ca is monic then $c_n = c^n$. It follows from Binet's formula that

$$\bar{\mathtt{Q}}_n = c^n \frac{f^{n+1} - (f')^{n+1}}{\sqrt{D}} = \frac{\bar{f}^{n+1} - (\bar{f}')^{n+1}}{c\sqrt{D}}, \quad \bar{f} := cf, \;\; \bar{f}' := cf'. \tag{4}$$

The set

$$\mathcal{B} = \{T^{d-1}\bar{\mathtt{Q}}_0, \dots, T\bar{\mathtt{Q}}_0, \bar{\mathtt{Q}}_0; T^{d-1}\bar{\mathtt{Q}}_1, \dots, T\bar{\mathtt{Q}}_1, \bar{\mathtt{Q}}_1; \dots\} \tag{5}$$

is a basis for A as an $\mathbb{F}_q$-vector space: as the degree map $\deg : \mathcal{B} \to \mathbb{N} = \{0, 1, \dots\}$ is a bijection. The order in which we have presented the elements of $\mathcal{B}$ corresponds to decreasing errors: for $0 \le l \le d-1$ and $n \ge 0$

$$\|T^l \bar{\mathtt{Q}}_n f\| = |T^l \bar{\mathtt{Q}}_n f - T^l \bar{\mathtt{Q}}_{n+1}| = q^{l-(n+1)d} < 1. \tag{6}$$

In particular, $(T^l \bar{\mathtt{Q}}_n)^{\perp} = T^l \bar{\mathtt{Q}}_n^{\perp}$, and the map

$$\mathcal{B} \longrightarrow q^{-\mathbb{N}}, \quad T^l \bar{\mathtt{Q}}_n \longmapsto \|T^l \bar{\mathtt{Q}}_n f\| \tag{7}$$

defines a bijection between $\mathcal{B}$ and the set of possible errors.

Write

$$\mathcal{B}(i) = \{T^{d-1}\bar{\mathtt{Q}}_i, \dots, \bar{\mathtt{Q}}_i\}$$

for the ith block of $\mathcal{B}$. Furthermore, for $0 \le \tilde{d} \le d-1$, denote

$$\mathcal{B}(i)_{\tilde{d}} = \{T^{\tilde{d}}\bar{\mathtt{Q}}_i, \dots, \bar{\mathtt{Q}}_i\}.$$

Lemma 1. *Let $l \in \{0, \dots, d-1\}$ and write*

$$d_l = d - 1 - l.$$

Then

$$\Lambda_{q^{-Nd-l}}(f) = \mathrm{span}_{\mathbb{F}_q}(\mathcal{B}(N)_{d_l}, \mathcal{B}(N+1), \dots).$$

Proof. First observe that

$$\mathrm{span}_{\mathbb{F}_q}(\mathcal{B}(N)_{d_l}, \mathcal{B}(N+1), \dots) \subset \Lambda_{q^{-Nd-l}}(f).$$

Indeed, for $T^{\tilde{d}}\bar{\mathsf{Q}}_{N+r}$ with $\tilde{d} \leq d-1$, and $\tilde{d} \leq d_l$ if $r = 0$, we have

$$\|T^{\tilde{d}}\bar{\mathsf{Q}}_{N+r} f\| \leq q^{d_l} \cdot q^{-(N+1)d} = q^{-Nd-l-1} < q^{-Nd-l}.$$

Moreover, by (6), $\Lambda_{q^{-Nd-l}}(f)$ contains no other elements of $\mathcal{B}$. In view of the bijection (7), no linear combination of the excluded basis elements could appear in $\Lambda_{q^{-Nd-l}}(f)$. This proves the equality in the statement of the Lemma. □

We recall the main definitions established in [2]. For each ε, $\Lambda_\varepsilon^{\mathrm{mon}}(f)$ is the subset of monic polynomials in $\Lambda_\varepsilon(f)$. We defined

$$\zeta_{f,\varepsilon}(n) := \sum_{\lambda \in \Lambda_\varepsilon^{\mathrm{mon}}(f) - \{0\}} \lambda^{-n}, \quad n \in \mathbb{N},$$

and

$$j_\varepsilon(f) = \frac{1}{\frac{1}{T^q - T} - J_\varepsilon(f)},$$

where

$$J_\varepsilon(f) = \frac{T^{q^2} - T}{(T^q - T)^{q+1}} \cdot \tilde{J}_\varepsilon(f), \quad \tilde{J}_\varepsilon(f) := \frac{\zeta_{f,\varepsilon}(q^2-1)}{\zeta_{f,\varepsilon}(q-1)^{q+1}}.$$

Then $j^{\mathrm{qt}}(f)$ is the set of limits of $j_\varepsilon(f)$ for $\varepsilon \to 0$. It follows from Lemma 1 that to calculate the values of $j^{\mathrm{qt}}(f)$, it suffices to consider the possible limits of $j_\varepsilon(f)$ formed from the spans of the initial truncations of $\mathcal{B}$.

Let $\varepsilon = q^{-Nd-l}$. As in [2], we decompose

$$\Lambda_\varepsilon^{\mathrm{mon}}(f) = \Lambda_\varepsilon^{\mathrm{bas}}(f) \bigsqcup \Lambda_\varepsilon^{\mathrm{non-bas}}(f)$$

where

$$\Lambda_\varepsilon^{\mathrm{bas}}(f) = \{\mathcal{B}(N)_{d_l}, \mathcal{B}(N+1), \dots\}$$

and

$$\Lambda_\varepsilon^{\mathrm{non-bas}}(f) = \Lambda_\varepsilon^{\mathrm{mon}}(f) - \left(\Lambda_\varepsilon^{\mathrm{bas}}(f) \cup \{0\}\right).$$

Any element $\lambda \in \Lambda_\varepsilon^{\mathrm{non-bas}}(f)$ may therefore be written in the form

$$\lambda = c_0(T)\bar{\mathsf{Q}}_N + c_1(T)\bar{\mathsf{Q}}_{N+1} + \cdots + c_m(T)\bar{\mathsf{Q}}_{N+m},$$

where the $c_i(T) \in A$ are polynomials that are not all zero and satisfy the following conditions

I. $c_m(T)$ is monic and if $\lambda = c_m(T)\bar{\mathsf{Q}}_{N+m}$, $c_m(T) \neq T^j$ for all $j \leq d-1$.
II. $\deg(c_i(T)) \leq d-1$, $i = 0, \dots, m$.
III. $\deg(c_0(T)) \leq d_l$.

In view of this characterization, sums over $\lambda \in \Lambda_\varepsilon^{\mathrm{non-bas}}(f)$ may be understood as sums indexed by tuples $c_0(T), \dots, c_m(T)$ subject to conditions I, II and III, and so we will abbreviate

$$\sum_{\substack{\text{Conditions}\\ \text{I,II,III}}} := \sum_{\Lambda_\varepsilon^{\mathrm{non-bas}}(f)}.$$

Using this notation, we may now express

$$\tilde{J}_\varepsilon(f) = \frac{\sum_{\lambda\in\Lambda_\varepsilon^{\mathrm{bas}}(f)} \lambda^{1-q^2} + \sum_{\substack{\text{Conditions}\\ \text{I,II,III}}} \left(\sum_{i=0}^m c_i(T)\bar{\mathbb{Q}}_{N+i}\right)^{1-q^2}}{\left(\sum_{\lambda\in\Lambda_\varepsilon^{\mathrm{bas}}(f)} \lambda^{1-q} + \sum_{\substack{\text{Conditions}\\ \text{I,II,III}}} \left(\sum_{i=0}^m c_i(T)\bar{\mathbb{Q}}_{N+i}\right)^{1-q}\right)^{q+1}}. \tag{8}$$

Write

$$\zeta_{T,l}(n) := 1 + T^{-n} + \cdots + T^{-nd_l}, \quad \zeta_T(n) := \zeta_{T,0}(n),$$

then define

$$H_l(n) := \zeta_{T,l}(n) + \sum_{\substack{\text{Conditions I,II,III}\\ c_0(T)\neq 0}} \left(\sum_{i=0}^m c_i(T)\bar{f}^i\right)^{-n}$$

and

$$H(n) := \frac{\zeta_T(n)}{\bar{f}^n - 1} + \sum_{\substack{\text{Conditions}\\ \text{I,II}}} \left(\sum_{i=1}^m c_i(T)\bar{f}^i\right)^{-n}.$$

Now let

$$\tilde{J}(f)_l := \frac{H_l(q^2-1) + H(q^2-1)}{\left(H_l(q-1) + H(q-1)\right)^{1+q}}.$$

Note that $\tilde{J}(f)_l$ converges: indeed both numerator and denominator have absolute value 1, and $|\tilde{J}(f)_l| = 1$. Finally, we write

$$J(f)_l := \frac{T^{q^2} - T}{(T^q - T)^{q+1}} \cdot \tilde{J}(f)_l.$$

Theorem 1. *Let f be a quadratic unit which is a solution of $X^2 - aX - b$ with $\deg a = d > 0$ and $\deg b = 0$. Then $j^{\mathrm{qt}}(f)$ has precisely $d = \deg a = \log_q|\sqrt{D}|$ values, and they are*

$$j^{\mathrm{qt}}(f) = \left\{ j(f)_l := \frac{1}{\frac{1}{T^q - T} - J(f)_l} \right\}_{l=0}^{d-1}.$$

Proof. The conjugate solution satisfies $f' = -bf^{-1}$ with $b \in \mathbb{F}_q^\times$, so f' is $\mathrm{GL}_2(A)$-equivalent to f, hence by modularity, $j^{\mathrm{qt}}(f') = j^{\mathrm{qt}}(f)$. Thus, in what follows, we will suppose f satisfies $|f| > 1$. As above, $\varepsilon = q^{-Nd-l}$. We begin by noting that

$$\sum_{\lambda\in\Lambda_\varepsilon^{\mathrm{bas}}(f)} \lambda^{-n} = \zeta_{T,l}(n)\bar{\mathbb{Q}}_N^{-n} + \zeta_T(n)\sum_{i=1}^\infty \bar{\mathbb{Q}}_{N+i}^{-n}.$$

Then if we define

$$H_{N,l}(n) := \zeta_{T,l}(n)\bar{\mathbb{Q}}_N^{-n} + \sum_{\substack{\text{Conditions I,II,III}\\ c_0(T)\neq 0}} \left(\sum_{i=0}^m c_i(T)\bar{\mathbb{Q}}_{N+i}\right)^{-n}$$

and

$$H_N(n) := \zeta_T(n) \sum_{i=1}^{\infty} \bar{\mathtt{Q}}_{N+i}^{-n} + \sum_{\substack{\text{Conditions} \\ \text{I,II}}} \left(\sum_{i=1}^{m} c_i(T) \bar{\mathtt{Q}}_{N+i} \right)^{-n},$$

we may re-write (8) as

$$\tilde{J}_\varepsilon(f) = \frac{H_{N,l}(q^2-1) + H_N(q^2-1)}{(H_{N,l}(q-1) + H_N(q-1))^{q+1}}. \tag{9}$$

Replace in (9) each $\bar{\mathtt{Q}}_{N+i}$ by $(\bar{f}^{N+i+1} - \bar{f}'^{N+i+1})/c\sqrt{D}$ using (4): equivalently (since the numerator and denominator of (9) have homogeneous degree $1-q^2$) we may omit the constant $1/c\sqrt{D}$ and simply replace $\bar{\mathtt{Q}}_{N+i}$ by $\bar{f}^{N+i+1} - \bar{f}'^{N+i+1}$. Then dividing out the numerator and denominator by $\bar{f}^{(1-q^2)(N+1)}$ yields

$$\tilde{J}_\varepsilon(f) = \frac{\hat{H}_{N,l}(q^2-1) + \hat{H}_N(q^2-1)}{\left(\hat{H}_{N,l}(q-1) + \hat{H}_N(q-1)\right)^{q+1}}, \tag{10}$$

where

$$\begin{aligned} \hat{H}_{N,l}(n) :=& \zeta_{T,l}(n) \left(1 - (\bar{f}'/\bar{f})^{N+1}\right)^{-n} \\ &+ \sum_{\substack{\text{Conditions I,II,III} \\ c_0(T) \neq 0}} \left(\sum_{i=0}^{m} c_i(T) \bar{f}^i \left(1 - (\bar{f}'/\bar{f})^{N+i+1}\right) \right)^{-n} \end{aligned}$$

and

$$\begin{aligned} \hat{H}_N(n) :=& \Bigg\{ \zeta_T(n) \sum_{i=1}^{\infty} \bar{f}^{-ni} \left(1 - (\bar{f}'/\bar{f})^{N+i+1}\right)^{-n} \\ &+ \sum_{\substack{\text{Conditions} \\ \text{I,II}}} \left(\sum_{i=1}^{m} c_i(T) \bar{f}^i \left(1 - (\bar{f}'/\bar{f})^{N+i+1}\right) \right)^{-n}. \end{aligned}$$

Since $|\bar{f}'/\bar{f}| < 1$,

$$1 - (\bar{f}'/\bar{f})^{N+i+1} \longrightarrow 1$$

as $N \to \infty$, uniformly in i, and it follows that

$$\lim_{N\to\infty} \hat{H}_{N,l}(n) = H_l(n), \quad \lim_{N\to\infty} \hat{H}_N(n) = H(n)$$

and therefore

$$\lim_{N\to\infty} \tilde{J}_{q^{-Nd-l}}(f) = \tilde{J}(f)_l.$$

□

Note 1. In the number field case, PARI GP experiments [3] performed on fundamental quadratic units θ indicate that $j^{\mathrm{qt}}(\theta)$ appears to have D values, where D is the corresponding fundamental discriminant.

3. Arbitrary Real Quadratics

Let $h \in k_\infty - k$ be an arbitrary quadratic. By the Dirichlet Unit Theorem [1], the group of units in the quadratic extension $k(h)$ is isomorphic to $\mathbb{F}_q^\times \times \mathbb{Z}$. Thus there exists a unit f such that $k(h) = k(f)$, i.e., h can be written in the form

$$h = \frac{x + yf}{z}, \quad x, y, z \in A, \tag{11}$$

where f satisfies $X^2 - aX - b = 0$, $d = \deg(a) > 0$ and $\deg(b) = 0$.

Note that for h and f as in (11) and ε satisfying $|z|\varepsilon < 1$, we have the following inclusions of $\mathbb{F}_q$-vector spaces

$$z\Lambda_{|y|^{-1}\varepsilon}(f) \subset \Lambda_\varepsilon(h) \subset y^{-1}\Lambda_{|z|\varepsilon}(f). \tag{12}$$

The first inclusion follows upon noting that if $z\lambda \in z\Lambda_{|y|^{-1}\varepsilon}(f)$ we have

$$\|(z\lambda)h\| = |(z\lambda)h - (y\lambda^\perp + \lambda x)| = |\lambda(x + yf) - (y\lambda^\perp + \lambda x)| = |y||\lambda f - \lambda^\perp| < \varepsilon.$$

The second inclusion follows from noting that if $\lambda \in \Lambda_\varepsilon(h)$, $y\lambda \in \Lambda_{|z|\varepsilon}(f)$, as

$$\|(y\lambda)f\| = |(y\lambda)f - (\lambda^\perp z - \lambda x)| = |\lambda(zh - x) - (\lambda^\perp z - \lambda x)| = |z||\lambda h - \lambda^\perp| < |z|\varepsilon.$$

Lemma 2. *The inclusion of $\mathbb{F}_q$-vector spaces*

$$z\Lambda_{|y|^{-1}\varepsilon}(f) \subset \Lambda_\varepsilon(h)$$

has index bounded by a constant which depends only on y and z.

Proof. By (12), it suffices to show that the induced inclusion

$$z\Lambda_{|y|^{-1}\varepsilon}(f) \subset y^{-1}\Lambda_{|z|\varepsilon}(f)$$

is of index bounded by a constant which only depends on y, z. For any $\delta < 1$, denote

$$\Lambda_\delta^2(f) := \{(\lambda, \lambda^\perp) \in A^2 |\ \lambda \in \Lambda_\delta(f)\}.$$

Note that the map $\lambda \mapsto (\lambda, \lambda^\perp)$ induces an isomorphism $\Lambda_\delta(f) \cong \Lambda_\delta^2(f)$ of $\mathbb{F}_q$-vector spaces: this in fact follows from Proposition 1 of [2]. In turn, we obtain an induced isomorphism of y^{-1}-rescalings

$$y^{-1}\Lambda_\delta(f) \cong y^{-1}\Lambda_\delta^2(f).$$

Taking $\delta = |z|\varepsilon$, the corresponding isomorphism takes $z\Lambda_{|y|^{-1}\varepsilon}(f)$ to $z\Lambda^2_{|y|^{-1}\varepsilon}(f)$. Thus

$$y^{-1}\Lambda_{|z|\varepsilon}(f)/z\Lambda_{|y|^{-1}\varepsilon}(f) \cong y^{-1}\Lambda^2_{|z|\varepsilon}(f)/z\Lambda^2_{|y|^{-1}\varepsilon}(f).$$

On the other hand, the natural map

$$y^{-1}\Lambda^2_{|z|\varepsilon}(f)/z\Lambda^2_{|y|^{-1}\varepsilon}(f) \hookrightarrow (y^{-1}A)^2/(zA)^2$$

is injective: for if $(\lambda_1, \lambda_1^\perp), (\lambda_2, \lambda_2^\perp) \in \Lambda^2_{|z|\varepsilon}(f)$ satisfy

$$y^{-1}(\lambda_1, \lambda_1^\perp) - y^{-1}(\lambda_2, \lambda_2^\perp) = z(\beta_1, \beta_2) \in (zA)^2,$$

then since $z(\beta_1, \beta_2) \in y^{-1}\Lambda^2_{|z|\varepsilon}(f)$ (being a difference of elements of the latter),

$$|(yz\beta_1)f - yz\beta_2| < |z|\varepsilon$$

which implies $(\beta_1, \beta_2) \in \Lambda^2_{|y|^{-1}\varepsilon}(f)$. Therefore the index of $z\Lambda_{|y|^{-1}\varepsilon}(f)$ in $y^{-1}\Lambda_{|z|\varepsilon}(f)$ is bounded by $\dim_{\mathbb{F}_q}((y^{-1}A)^2/(zA)^2)$. This proves the Lemma. □

Theorem 2. *Let $h \in k_\infty - k$ be quadratic. Then $\#j^{\mathrm{qt}}(h) < \infty$.*

Proof. By Lemma 2, for each ε, we may find $\lambda_{1,\varepsilon} = 0, \ldots, \lambda_{r,\varepsilon} \in \Lambda_\varepsilon(h)$, $r < M = M(y,z)$, such that

$$\Lambda_\varepsilon(h) = \big[z\Lambda_{|y|^{-1}\varepsilon}(f) + \lambda_{1,\varepsilon}\big] \bigsqcup \cdots \bigsqcup \big[z\Lambda_{|y|^{-1}\varepsilon}(f) + \lambda_{r,\varepsilon}\big]. \tag{13}$$

Thus, we have

$$\begin{aligned}\tilde{J}_\varepsilon(h) = \frac{\zeta_{h,\varepsilon}(q^2-1)}{\zeta_{h,\varepsilon}(q-1)^{q+1}} &= -\frac{\sum_{\lambda\in\Lambda_\varepsilon(h)-0}\lambda^{1-q^2}}{\Big(\sum_{\lambda\in\Lambda_\varepsilon(h)-0}\lambda^{1-q}\Big)^{q+1}} \\ &= -\frac{\sum_{i=1}^r\sum_{\lambda\in\Lambda_{|y|^{-1}\varepsilon}(f)}(\lambda+\lambda_{i,\varepsilon}/z)^{1-q^2}}{\Big(\sum_{i=1}^r\sum_{\lambda\in\Lambda_{|y|^{-1}\varepsilon}(f)}(\lambda+\lambda_{i,\varepsilon}/z)^{1-q}\Big)^{1+q}}.\end{aligned}$$

By (12), for $i \neq 1$,

$$\lambda_{i,\varepsilon} \in y^{-1}\Lambda_{|z|\varepsilon}(f) - z\Lambda_{|y|^{-1}\varepsilon}(f), \tag{14}$$

so we may write

$$\lambda_{i,\varepsilon} = y^{-1}\sum_{j=m}^{m'} c_{ij,\varepsilon}(T)\bar{\mathtt{Q}}_i, \tag{15}$$

where $\deg(c_{ij}(T)) \leq d-1$ and m is the smallest index so that $\bar{\mathtt{Q}}_m \in \Lambda_{|z|\varepsilon}(f)$. Note that by (14) and Lemma 2, the difference $m'-m$ has a uniform bound which is independent of ε. In particular, the set of all possible coefficients $c_{ij,\varepsilon}(T)$ is finite in number.

Consider a sequence $\{q^{-Nd-l}\}_{N>1}$ of values for ε giving a limiting value for $j^{\mathrm{qt}}(f)$ as described in Theorem 1: recall that the limit is calculated by using Binet's formula to replace occurrences of $\bar{\mathtt{Q}}_{N+i}$ by $\bar{f}^{N+i+1}$ in $\tilde{J}_\varepsilon(f)$ and dividing out by $\bar{f}^{(1-q^2)(N+1)}$. We apply exactly the same process to $\tilde{J}_\varepsilon(h)$: and in view of (15) we obtain approximations

$$z^{-1}\frac{\lambda_{i,\varepsilon}}{\bar{f}^{N+1}} \sim (yz)^{-1}\sum_{j=m}^{m'} c_{ij,\varepsilon}(T)\bar{f}^{j-N}, \quad m, m' \geq 0 \tag{16}$$

where $\deg c_{ij,\varepsilon}(T) \leq d-1$. Therefore, the set of possible limits of the $z^{-1}\frac{\lambda_{i,\varepsilon}}{\bar{f}^{N+1}}$ as $N \to \infty$ is contained in the finite set

$$\left\{ (yz)^{-1}\sum_{j=-n}^{n'} c_j(T)\bar{f}^j \;\middle|\; \deg(c_j(T)) \leq d-1 \right\}$$

where $n'+n = m'-m$. Thus, within the family $\{\varepsilon = q^{-Nd-l}\}_{N>1}$, there are only finitely many possible limits of sub-sequences of $\{\tilde{J}_\varepsilon(h)\}$ giving rise to elements of $j^{\mathrm{qt}}(h)$. This proves the Theorem. □

References

[1] Cohn, P.M., *Algebraic Numbers and Algebraic Functions.* Chapman and Hall/CRC, London, 1991.

[2] Demangos, L & Gendron, T.M., Quantum j-Invariant in positive characteristic I: definitions and convergence.

[3] Castaño Bernard, C. & Gendron, T.M., Modular invariant of quantum tori. *Proc. Lond. Math. Soc.* **109** (2014), Issue 4, 1014–1049.

Instituto de Matemáticas – Unidad Cuernavaca, Universidad Nacional Autonoma de México, Av. Universidad S/N, C.P. 62210 Cuernavaca, Morelos, México
Current address: Department of Mathematical Sciences (Mathematics Division), University of Stellenbosch, Private Bag X1, Matieland 7602, South Africa
E-mail address: l.demangos@gmail.com

Instituto de Matemáticas – Unidad Cuernavaca, Universidad Nacional Autonoma de México, Av. Universidad S/N, C.P. 62210 Cuernavaca, Morelos, México
E-mail address: tim@matcuer.unam.mx